



National EMS Quality Alliance

Trauma-04 Measure Package

Disclaimer

EMS Performance Measures (Measures) and related data specifications developed by the National EMS Quality Alliance (NEMSQA) are intended to facilitate quality improvement activities by EMS professionals.

These measures are intended to assist EMS professionals in enhancing quality of care. These Measures are not clinical guidelines and do not establish a standard of medical care and have not been tested for all potential applications. NEMSQA encourages testing and evaluation of its Measures.

Measures are subject to review and may be revised or rescinded at any time by NEMSQA. The measures may not be altered without prior written approval from NEMSQA. The measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes (e.g., use by health care providers in connection with their practices). Commercial use is defined as the sale, license, or distribution of the measures for commercial gain, or incorporation of the measures into a product or service that is sold, licensed, or distributed for commercial gain. Commercial uses of the measures require a license agreement between the user and NEMSQA. Neither NEMSQA nor its members shall be responsible for any use of the measures.

THESE MEASURES AND SPECIFICATIONS ARE PROVIDED “AS IS” WITHOUT WARRANTY OF ANY KIND.

©2019 National EMS Quality Alliance. All rights reserved.

Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary coding sets should obtain all necessary licenses from the owners of these code sets. NEMSQA and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications. ICD-10 copyright 2019 International Health Terminology Standards Development Organization.

CPT ® is a registered trademark of the American Medical Association and is copyright 2019. CPT® codes contained in the Measure specifications are copyright 2004-2019 American Medical Association

Trauma-04: Trauma Patients Transported to the Trauma Center

Trauma-04 was designed using CDC guidelines for Field Triage of Trauma Patients. Along with the CDC guidelines, published literature clearly supports this measure, as patients who receive appropriate trauma care often have better outcomes. As this measure was being re-specified, the TEP discussed many concerns with the pragmatic implementation of this measure – including the availability of trauma centers in rural communities and whether the measure should focus on transporting patients to the appropriate level of trauma center. While there are many possible variations and stratifications of this measure, the TEP ultimately decided to stay with the intent of the original EMS Compass candidate measure, which is measuring if patients with trauma are being transported to a trauma center.

During the re-specification process, the TEP closely reviewed the CDC Guidelines for Field Triage, which were used to build the denominator for this measure. Originally, in the EMS Compass candidate measures, Step 1, 2, and 3 criteria were part of the denominator inclusion criteria. But, after reviewing the guidelines again, the TEP decided to remove Step 3 from the denominator for the re-specified measure, as Steps 1 and 2 identify the most seriously injured patients. The experts determined that limiting the denominator to **Steps 1 and 2** will satisfy the intent of the measure without running the risk of over transporting patients to trauma centers who may not need the care of such a facility.

As stated above, much discussion was had about the numerator of Trauma-04. While everyone on the TEP agreed that transporting a trauma center is the best course of treatment for certain patients, it was noted that trauma centers are not always available or well-defined. The TEP discussed many different options for the numerator for Trauma-04, including transporting patient with trauma to the nearest hospital and transporting patients to a specific level of trauma center. However, the final measure has been specified to require the patient to be transported to a trauma center.

The TEP understands concerns about limited access to trauma centers in certain communities and concerns that some hospitals labeled as trauma centers may not be as equipped as others. But, as measurement is used to drive change, the TEP hopes data collected from Trauma-04 will help drive change in the availability and standards of trauma centers across the nation.

Trauma-04: Trauma Patients Transported to a Trauma Center


Measure Score Interpretation: For this measure, a higher score indicates better quality.

Measure Description	
Percentage of EMS responses originating from a 911 request for patients who meet CDC criteria for trauma and are transported to a trauma center.	
Measure Components	
Numerator Statement	EMS transports originating from a 911 request for patients transported to a trauma center.
Denominator Statement	All EMS transports originating from a 911 request for patients who meet 2011 CDC Step 1 or 2 criteria for trauma.
Denominator Exclusions	None
Denominator Exceptions	None
Supporting Guidance & Other Evidence	<p>The following evidence statements are quoted verbatim from the referenced treatment protocol:</p> <p>Centers for Disease Control and Prevention: Guidelines for Field Triage of Injured Patients:¹</p>
Measure Importance	

Rationale	<p>Victims of severe violent injuries involving trauma not only see a slight improvement in survival rates if they receive treatment in a trauma center, but they also benefit from less complications and shorter lengths of stay.ⁱⁱ</p> <p>A study on the effect of trauma center care on mortality published in 2006 found that one-year survival rates among patients with traumatic injuries were significantly higher when patients received care in trauma centers as opposed to non-trauma centers. This same study showed a 25% decrease in mortality for severely injured adult patients who received care at Level I trauma centers.ⁱⁱⁱ</p>
Measure Designation	
Measure purpose	<ul style="list-style-type: none"> • <input checked="" type="checkbox"/> Quality Improvement • <input checked="" type="checkbox"/> Accountability • <input type="checkbox"/> MOC
Type of measure	<ul style="list-style-type: none"> • <input checked="" type="checkbox"/> Process • <input type="checkbox"/> Outcome • <input type="checkbox"/> Structure • <input type="checkbox"/> Efficiency
National Quality Strategy/Priority/CMS Measure Domain	<ul style="list-style-type: none"> • <input checked="" type="checkbox"/> Clinical Process-Effectiveness • <input type="checkbox"/> Patient Safety • <input type="checkbox"/> Patient Experience • <input type="checkbox"/> Care Coordination • <input type="checkbox"/> Efficiency: Overuse • <input type="checkbox"/> Efficiency: Cost • <input type="checkbox"/> Population & Community Health
CMS Meaningful Measure Domain	<ul style="list-style-type: none"> • <input type="checkbox"/> Medication Management • <input type="checkbox"/> Admissions and Readmissions to Hospitals • <input type="checkbox"/> Transfer of Health Information and Interoperability • <input type="checkbox"/> Preventative Care • <input type="checkbox"/> Management of Chronic Conditions • <input type="checkbox"/> Prevention, Treatment, and Management of Mental Health • <input type="checkbox"/> Prevention and Treatment of Opioid and Substance • <input type="checkbox"/> Risk Adjusted Mortality • <input type="checkbox"/> Equity of Care • <input type="checkbox"/> Community Engagement • <input type="checkbox"/> Appropriate Use of Healthcare • <input type="checkbox"/> Patient-focused Episode of Care • <input type="checkbox"/> Risk-Adjusted Total Cost of Care • <input type="checkbox"/> Healthcare-associated infections • <input type="checkbox"/> Preventable Healthcare Harm

	<ul style="list-style-type: none"> • <input checked="" type="checkbox"/> Care is Personalized and Aligned with Patient's Goals • <input type="checkbox"/> End of Life Care according to Preferences • <input type="checkbox"/> Patient's Experience of Care • <input type="checkbox"/> Patient Reported Functional Outcomes
Level of measurement	<ul style="list-style-type: none"> • <input checked="" type="checkbox"/> Individual EMS Professional • <input checked="" type="checkbox"/> EMS Agency
Care setting	<ul style="list-style-type: none"> • <input checked="" type="checkbox"/> Pre-Hospital Care
Data source	<ul style="list-style-type: none"> • <input checked="" type="checkbox"/> Electronic Patient Care Record (eCPR) data • <input type="checkbox"/> Administrative Data/Claims (inpatient, outpatient or multiple-source claims) • <input checked="" type="checkbox"/> Paper medical record/Chart abstracted • <input checked="" type="checkbox"/> Registry

Clinical Quality Measure Flow for Trauma-04 Trauma Patients Transported to a Trauma Center



ⁱ Sasser, S.M., Hunt, R.C., Faul, M., Sugerman, D., Pearson, W.S., Dulski, T., Wald, M.M., Jurkovich, G.J., Newgard, C.D., Lerner, E.B., Cooper, A., Wang, S.C., Henry, M.C., Salomone, J.P., Galli, R.L. (2011) Guidelines for Field Triage of Injured Patients: Recommendations of the National Expert Panel on Field Triage, 2011. *MMWR Morb Mortal Wkly Rep*; 61(RR01), 1-20.

ⁱⁱ Baez, A.A., Lane, P.L., Sorondo, B., Nituica, C. (2006) Receiving Care Facility and Outcome Differences for Victims of Severe Violent injuries, *Prehospital Emergency Care*, 10:2, 220-223.

ⁱⁱⁱ MacKenzie, E.J., Rivara, F.P., Jurkovich, G.J., Nathens, A.B., Frey, K.P., Egleston, B.L., Salkever, D.S., Scharfstein, D.O., (2006) A National Evaluation of the Effect of Trauma-Center Care on Mortality, *The New England Journal of Medicine*, 354;4, 366-378.